

## Chocolate Quality Testing

Chocolate appears in many forms and as an ingredient in many confectionery and non-confectionery prod-



*Chocolate has many forms and is an ingredient in many products.*

Photo © iStockphoto.com/FotografiaBasica

ucts. The December 2010 issue of *Food Technology* contains an overview of the chocolate confectionery market. This article focuses on the production and quality testing of chocolate.

### Chocolate Production

Cacao trees produce pods throughout the year, but large harvests occur twice annually.

Workers harvest the pods by hand, hack open the shells, scoop out the pulp and seeds, and discard the husks. The pulp and seeds are placed into boxes or piles, covered with

banana leaves, and left to ferment for up to 9 days, during which flavor and aroma compounds are formed. The seeds are then dried in the sun or in dryers for about a week, packed into burlap bags, and shipped to manufacturing plants. There they are mechanically cleaned to remove dried pulp and debris, sorted, and roasted to release the flavor and aroma compounds. The roasting also deepens the brown color of the beans.

The beans are then lightly crushed, and the husks are blown away in a process called winnowing, leaving dark chips called nibs whose cells contain about 53% fat (cocoa butter) and 47% solids. The nibs are then crushed or ground in mills that generate enough friction and heat to liquefy the nibs into a thick paste called chocolate liquor. Some of the chocolate liquor is placed in a hydraulic press that squeezes out the cocoa butter, and the remaining solid cocoa is pulverized into cocoa powder.

The chocolate liquor can also be poured into molds to harden into unsweetened chocolate, which contains 100% chocolate liquor. To produce dark, semisweet, and bittersweet chocolates, nibs are ground with sugar to reduce the particle size. The resulting paste is then kneaded in a process called conching to further smooth the texture. Aerating during conching creates a well-rounded flavor. The conched chocolate is then tempered by repeatedly

cooling and heating it to provide a glossy sheen and ensure that the chocolate will melt properly. The tempered chocolate can be molded or poured over flavored centers (enrobing) or used as a filling in other confections.

To make baking chocolate and candy bars, chocolate liquor is further processed with sugar and other ingredients. Chocolates can vary in amounts of cocoa particles, cocoa butter, sugar, and other ingredients such as emulsifiers and dairy products.

The U.S. Food and Drug Administration (FDA) has established standards of identity for cacao-derived products. Milk chocolate contains cream or other dairy products and sugar and must contain at least 10% chocolate liquor. Dark, bitter-sweet, or semisweet chocolate must contain at least 35% chocolate liquor but generally contains 50%–55% with the rest being sugar. High-percentage chocolates contain even more chocolate liquor and consequently more cocoa particles and cocoa butter than regular chocolate. Couverture, used for coating, contains more cocoa butter so that it is more free-flowing when melted. White chocolate contains no chocolate liquor but instead consists of cocoa butter, sugar, dairy products, and flavorings; it must contain at least 20% cocoa butter and no more than 55% sugar.

Title 21, Part 163, of the *Code of Federal Regulations* details





final chocolate. Flavor testing is absolutely critical in roasting and helps determine the degree of roast, Stauffer said. Every operator is trained in sensory testing, and they look for off-notes right after roasting and after chocolate liquor grinding. Sensory panels check all final chocolate products.

Another final test of deposited products is determining the count per pound. Chocolate chips for baking are typically sized for grocery store sales as 1,000 count per pound, but the company also provides counts of 2,000, 4,000, and 10,000, depending on what its customers in the baking, confectionery, dairy, and functional foods industries want.

#### **Challenges Ahead & Research Needed**

Stauffer said that she would like to see rapid methods for determining fat and moisture content and automatic titrations to determine fat quality. Current

instruments require extensive calibration to maintain, she said, so it would be great to have rapid methods for fat and moisture determinations as well as rapid microbiological methods. Testing for yeast and mold takes five days, but the product needs to be out the door faster, she said. Blommer outsources microbiological and chemical testing to independent laboratories. Most labs provide a 10-day turnaround for chemical testing, but two days is preferable. Sugar testing can take up to eight days. The alternative is to have testing equipment in house and have technicians do the testing, she added, but that's an additional expense.

Stauffer pointed out that there is a need for standardization of methods for testing chocolate. Temper, she said, is always difficult to measure quickly and constantly, viscosity measurements vary among technicians and plants, and the equipment



*Chocolate is created from the pulp and seeds of pods from cacao trees.*

Photo © iStockphoto.com/JackJelly

## Chocolate Quality Testing continued...

### Organizations Related to Chocolate

#### **National Confectioners Association** ([www.candyusa.com](http://www.candyusa.com)).

Founded in 1884, NCA is a trade association of confectionery companies. Its Chocolate Council was founded in 2008 by member companies that process cocoa and make chocolate. It focuses on educating the public about chocolate. NCA's Sweets & Snacks Expo will be held May 8–10, 2012, in Chicago, Ill.

**American Association of Candy Technologists** ([www.aactcandy.org](http://www.aactcandy.org)). AACT is a professional organization of individual technologists, operations personnel, educators, students, business staff, and others dedicated to the advancement of the confectionery industry. AACT's 2012 National AACT Technical Seminar will be held October 1–3, 2012, in Lincolnshire, Ill.

**PMCA** ([www.pmca.com](http://www.pmca.com)). Founded in 1907 as the Pennsylvania Manufacturing Confectioners' Association, PMCA is a trade association of confectionery manufacturers and companies in related industries. Its 66th Annual Production Conference will be held April 30–May 2, 2012, in Hershey, Pa. PMCA's Research Committee directs basic and applied research in the science of chocolate and confectionery at several universities. PMCA also presents short courses in confectionery technology.

#### **Retail Confectioners International** ([www.retailconfectioners.org](http://www.retailconfectioners.org)).

Founded in 1917, RCI is a trade association of chocolate makers, confectioners, and industry suppliers. It provides a forum for confectioners to meet, network, share ideas, solve mutual problems, and develop their candy-making and entrepreneurial skills. It presents short courses for retailers, including "Chocolate Boot Camp." RCI's 2012 Annual Convention & Industry Exposition will be held June 11–15, 2012, in San Francisco, Calif.

**International Confectionery Association** ([www.international-confectionery.com](http://www.international-confectionery.com)). ICA, formerly the International Office of Cocoa, Chocolate and Sugar Confectionery, is an organization of chocolate and confectionery manufacturers and suppliers of ingredients, machinery, packaging materials, and services to the industry. It facilitates, coordinates, and communicates international scientific, regulatory, and public affairs information while promoting and representing the interests of the global confectionery industry.

**World Cocoa Foundation** ([www.worldcocoafoundation.org](http://www.worldcocoafoundation.org)). WCF is an organization of manufacturers, processors, supply chain managers, ports, and allied industries and trade associations that works to promote sustainable cocoa production and improve the well-being of cocoa farmers through economic and social development and environmental stewardship.

can be difficult to calibrate. Particle-size measuring systems can also show variation. The main issue, she said, is that there is not always a standard method for the chocolate industry to follow. Companies around the world have their own ways of measuring and their own types of equipment. As a result, Blommer uses four different fat analyzers.

Ziegler said that finding a way to visualize the fat crystal network in chocolate, including the size and distribution, could provide more information about bloom stability—resistance to oil and moisture migration—and lead to a way to predict bloom in chocolate and chocolate-filled confections. He said that Penn State researchers are studying chocolate structure and its effect on sensory quality. There is also a correlation between mouthfeel and flavor, and changing the viscosity has dramatic effects on mouthfeel. Besides particle size, there is also interest in studying the shape of all the particles in chocolate, not only the cocoa but also the sugar and milk solids, he said. If products with the same particle size don't behave quite the same, some other factor must be involved. **FT**



**Neil H. Mermelstein**, a Fellow of IFT, is Editor Emeritus of Food Technology  
• [nhmermelstein@comcast.net](mailto:nhmermelstein@comcast.net)